arXiv:1201.5770v1 [hep-ph] 27 Jan 2012

The strong running coupling at τ and Z_0 mass scales from lattice QCD

(ETM Collaboration) B. Blossier,[1] Ph. Boucaud,[1] M. Brinet,[2] F. De Soto,[3]
X. Du,[2] V. Morenas,[4] O. Pène,[5] K. Petrov,[5] and J. Rodríguez-Quintero[6]

[1] *Laboratoire de Physique Théorique, Université de Paris XI; Bâtiment 210, 91405 Orsay Cedex; France*
[2] *Laboratoire de Physique Subatomique et de Cosmologie,*
CNRS/IN2P3/UJF; 53, avenue des Martyrs, 38026 Grenoble, France
[3] *Dpto. Sistemas Físicos, Químicos y Naturales, Univ. Pablo de Olavide, 41013 Sevilla, Spain*
[4] *Laboratoire de Physique Corpusculaire, Université Blaise Pascal, CNRS/IN2P3 63177 Aubière Cedex, France*
[5] *Laboratoire de Physique Théorique, Université de Paris XI; Bâtiment 211, 91405 Orsay Cedex; France*
[6] *Dpto. Física Aplicada, Fac. Ciencias Experimentales; Universidad de Huelva, 21071 Huelva; Spain.*

This letter reports on the first computation, from data obtained in lattice QCD with u, d, s and c quarks in the sea, of the running strong coupling via the ghost-gluon coupling renormalized in the MOM Taylor scheme. We provide with estimates of $\alpha_{\overline{\rm MS}}(m_\tau^2)$ and $\alpha_{\overline{\rm MS}}(m_Z^2)$ (for which the inclusion of the dynamical charm quark makes the running much safer) in very good agreement with experimental results.

PACS numbers: 12.38.Aw, 12.38.Lg

INTRODUCTION

The confrontation of QCD, the theory for the strong interactions, with experiments requires a few inputs: one mass parameter for each quark species and the only surviving parameter in the limit of massless quarks, namely $\Lambda_{\rm QCD}$, the energy scale used as the typical boundary condition for the integration of the Renormalization Group equation for the strong coupling constant, α_S. Thus, the value of the renormalized strong coupling at any scale, or equivalently $\Lambda_{\rm QCD}$, has to be fitted to allow the QCD phenomenology to account successfully for experiments (see ref. [1] for a description of many precision measurements of α_S from different proccesses and at different energy scales). The QCD running coupling can be also obtained from lattice computations, where the lattice spacing replaces $\Lambda_{\rm QCD}$ as a free parameter to be adjusted from experimental numbers. This means that one can apply QCD implemented via a lattice-regularized action to convert the physical observation used for the lattice spacing calibration, as for instance a mass or a decay constant, into $\Lambda_{\rm QCD}$. A review of most of the procedures recently implemented to determine the strong coupling from the lattice can be found in ref. [2] (see also ref. [3], where we quoted many of the different methods proposed in the last few years).

The present "*world average*" for the strong coupling determinations [4], usually referred to at the scale of the Z^0 mass, as discussed in ref. [1], is dominated by the lattice determination included in the average [5]. Provided the importance of a precise and proper knowledge of the strong coupling for the LHC cross sections studies and its exploration of new physics, independent alternative lattice determinations are strongly required. The latter is specially true when different lattice actions and procedures are applied, to gain thus the best possible control for any source of systematic uncertainty. Furthermore, the current lattice results have been obtained by means of simulations including only two degenerate up and down sea quarks (N$_f$=2) or, as much (in particular that in [5]), also including one more "tuned" to the strange quark (N$_f$=2+1). Now, the European Twisted Mass (ETM) collaboration have started a wide-ranging program of lattice QCD calculation with two light degenerate twisted-mass flavours [6, 7] and a heavy doublet for the strange and charm dynamical quarks (N$_f$=2+1+1) [8, 9]. Within this ETM program, we have applied the method to study the running of the strong coupling, and so evaluate $\Lambda_{\rm QCD}$, grounded on the lattice determination of the ghost-gluon coupling in the so-called MOM Taylor renormalization scheme [10, 11]. We are publishing the results of this study in two papers: a methodological one [3], where the procedure is dicussed and described in detail and some results are presented, and this short letter aimed to update and emphasize the phenomenologically relevant results. In particular, as far as the lattice gauge fields with 2+1+1 dynamical flavours which we are exploiting provide with a very realistic simulation of QCD at the energy scales for the τ physics, we are presenting here the estimate for the coupling at the τ-mass scale and directly comparing with the one obtained from τ decays. It should be noted that including the dynamical charm quark makes also safer the running up to the Z^0-mass scale.

THE STRONG COUPLING IN TAYLOR SCHEME

The starting point for the analysis of this letter shall be the strong running coupling renormalized in the MOM-like Taylor scheme,

$$\alpha_T(\mu^2) \equiv \frac{g_T^2(\mu^2)}{4\pi} = \lim_{\Lambda\to\infty} \frac{g_0^2(\Lambda^2)}{4\pi} G(\mu^2,\Lambda^2) F^2(\mu^2,\Lambda^2) \ , \tag{1}$$

obtained from lattice QCD simulations; where F and G stand for the form factors of the two-point ghost and gluon Green functions (dressing functions). The procedure to compute the coupling defined by (1), and from it to extract an estimate of $\Lambda_{\overline{\rm MS}}$, is described in very detail in refs. [10, 11] and we recently applied this in ref. [3] to compute $\Lambda_{\overline{\rm MS}}$ from N_f=2+1+1 gauge configurations for several bare couplings, light twisted masses and volumes. The prescriptions applied for the appropriate elimination of discretization artefacts, as the so-called $H(4)$-extrapolation procedure to cure the artefacts which are due to the breaking of the rotational invariance on the lattice [12], needed to be provided with reliable and exploitable results were also carefully shown and explained in ref. [3]. After this, we are left with the lattice estimates of the Taylor coupling computed over a large range of momenta that, as can be seen in fig. 1, can be described above around 4 GeV by the following OPE formula:

$$\alpha_T(\mu^2) \ = \ \alpha_T^{\rm pert}(\mu^2) \ \left(1 + \frac{9}{\mu^2} \ R\left(\alpha_T^{\rm pert}(\mu^2), \alpha_T^{\rm pert}(q_0^2)\right) \right. \\ \left. \times \ \left(\frac{\alpha_T^{\rm pert}(\mu^2)}{\alpha_T^{\rm pert}(q_0^2)} \right)^{1-\gamma_0^{A^2}/\beta_0} \ \frac{g_T^2(q_0^2)\langle A^2\rangle_{R,q_0^2}}{4(N_C^2-1)} \right) , \quad (2)$$

derived from the OPE description of ghost and gluon dressing functions [11], where $\gamma_0^{A^2}$ can be taken from [13, 14] to give for $N_f = 4$, $1-\gamma_0^{A^2}/\beta_0 = 27/100$, and, applying the same method outlined in the appendix of ref. [11], one can take advantage of the $\mathcal{O}(\alpha^4)$-computations for the Wilson coefficients in ref. [14], and obtains $R(\alpha,\alpha_0)$ for $q_0 = 10$ GeV (see Eq.(6) of [3]). The purely perturbative running in Eq. (2) is given up to four-loops by integration of the β-function [4], where its coefficients are taken to be defined in Taylor-scheme [10, 15]. The inclusion of the non-perturbative OPE correction is unavoidable to properly describe the running of the coupling over the accesible momentum range, at least above around 4 GeV. This however allows to fit both $g^2\langle A^2\rangle$ and Λ_T, the $\Lambda_{\rm QCD}$ parameter in Taylor scheme, through the comparison of the prediction given by Eq. (2) and the lattice estimate of Taylor coupling. The best-fit of Eq. (2) to the lattice data published in ref. [3] can be seen in the plot of Fig. 1 (dotted line), while the best-fitted parameters should be read in Tab. I. In this letter, we also update the analysis by including an "*ad-hoc*" correction to account for higher power corrections (suggested by the detailed analysis of ref. [3]) that allows to extend the fitting window down to $p \simeq 1.7$ GeV. Furthermore, in addition to the lattice ensembles of gauge configurations described in ref. [3], we study 60 more at $\beta = 2.1$ ($a\mu_l = 0.002$) and three new ensembles of 50 configurations at $\beta = 1.9$ and $a\mu_l = 0.003, 0.004, 0.005$ to peform a chiral extrapolation for the ratios of lattice spacings and get: $a(2.1, 0.002)/a(1.9, 0) = 0.685(21)$. Thus, we have finally converted our estimates to physical units with the previous ratio and the lattice size in the chiral limit at $\beta = 1.9$ [8]: 0.08612(42) fm. .



FIG. 1: The strong running coupling in Taylor scheme defined by (1) obtained over a large momentum range from lattice QCD, as described in ref. [3]. The dotted line stands here for the best-fit with Eq. (2), while the solid one includes a higher-order power correction effectively behaving as $\sim 1/p^6$.

THE WILSON OPE COEFFICIENT AND THE HIGHER-POWER CORRECTIONS

The OPE prediction for α_T given by Eq. (2) is dominated by the first correction introduced by the non-vanishing dimension-two landau-gauge gluon condensate [16–21], where the Wilson coefficient is applied at the $\mathcal{O}(\alpha^4)$-order. In the previous technical paper [3], we provided with a strong indication that the OPE analysis is indeed in order: it was clearly shown that the lattice data could be only explained by including non-perturbative contributions and that the Wilson coefficient for the Landau-gauge gluon condensate was needed to describe the behaviour of data above $p \simeq 4$ GeV.

In Fig. 2, the impact of higher-power corrections is analized. The plot shows the departure of the lattice data for the Taylor coupling from the prediction given by Eq. (2), plotted in terms of the momentum, with logarithmic scales for both axes. The data seem to indicate that the next-to-leading non-perturbative correction is highly dominated by an $1/p^6$ term. This might suggest that the $1/p^4$ OPE contributions are negligible when compared with the $1/p^6$ ones or that the product of the leading $1/p^4$ terms and the involved Wilson coefficients leave with an effective $1/p^6$ behaviour. Anyhow, this implies that we can effectively describe the lattice data for the Taylor coupling with

$$\alpha_T^d(p^2) \ = \ \alpha_T(p^2) \ + \ \frac{d}{p^6} \ , \quad (3)$$

where d is a free parameter to be fitted, which we do not attribute to any particular physical meaning, for all



FIG. 2: The departure of lattice data from the leading non-perturbative OPE prediction for the running coupling plotted in logarithmic scales, in terms of the momentum manifestly shows a next-to-leading $1/p^6$ behaviour; the dotted red line stands for the momentum scale, $p \simeq 1.7$ GeV, below which the lattice data do not follow any more the $1/p^6$ behaviour.

momenta below $p \simeq 1.7$ GeV. This can be seen in the plot of Fig. 1, where the solid black line shows the best-fit of Eq. (3) to the lattice data, with the best-fit parameters that can be found in Tab. I. This table clearly shows that the values for the best-fit parameters obtained with a formula including the effective higher power correction applied for momenta above $p \simeq 1.7$ GeV are compatible with the ones obtained without higher power corrections and for momenta above $p \simeq 4$ GeV.

THE STRONG COUPLING IN $\overline{\text{MS}}$ SCHEME

To obtain the strong $\overline{\text{MS}}$ running coupling from the previously estimated values of Λ_T is rather immediate, as the scale-independent Λ_{QCD}-parameters in both Taylor and $\overline{\text{MS}}$ schemes are related through [11]

$$\frac{\Lambda_{\overline{\text{MS}}}}{\Lambda_T} = e^{-\dfrac{507-40N_f}{792-48N_f}} = 0.560832 \ . \tag{4}$$

Then, the $\overline{\text{MS}}$ running coupling can be again obtained by the integration of β-function, with the coefficients now in $\overline{\text{MS}}$-scheme for $\text{N}_f = 4$. Thus, we can apply the estimates of $\Lambda_{\overline{\text{MS}}}$, that can be found in Tab. I, to run the coupling down to the scale of τ mass, below the bottom quark mass threshold, and compare the result with the estimate from τ decays [1], $\alpha(m_\tau^2) = 0.334(14)$. This can be seen graphically in the plot of Fig. 3, while the results will be shown in the following.

	$\Lambda_{\overline{\text{MS}}}^{N_f=4}$ (MeV)	$g^2\langle A^2\rangle$ (GeV^2)	$d^{1/6}$ (GeV)
Eq. (2) [3]	316(13)	4.5(4)	
Eq. (3)	324(17)	3.8(1.0)	1.72(3)

TABLE I: The parameters for the best-fit of Eq. (2) obtained in ref. [3] to lattice data (first row) and the same with Eq. (3) (second row). The conversion to $\overline{\text{MS}}$ scheme for Λ_{QCD} is done by applying Eq. (4). The renormalization point for the gluon condensate is fixed at $\mu = 10$ GeV. We quote statistical errors obtained by applying the Jacknkife's method.

The running of $\alpha_{\overline{\text{MS}}}$ with the two estimates of $\Lambda_{\overline{\text{MS}}}$ from Tab. I, and the 1-σ error propagation, will produce the two following results at the τ-mass scale: $\alpha_{\overline{\text{MS}}}(m_\tau^2) = 0.337(8)$ and $\alpha_{\overline{\text{MS}}}(m_\tau^2) = 0.342(10)$. If we combine both and conservatively add the errors in quadrature, we will be left with the following estimate,

$$\alpha_{\overline{\text{MS}}}(m_\tau^2) \ = \ 0.339(13) \ , \tag{5}$$

in very good agreement with the one from τ decays.



FIG. 3: The strong $\overline{\text{MS}}$ coupling running for 4 quark flavours and for $\Lambda_{\overline{\text{MS}}} = 316$ MeV (black) and $\Lambda_{\overline{\text{MS}}} = 324$ MeV (blue) below the bottom mass threshold. The dotted lines represent the one-σ statistical deviations. The red point stand for the value of $\alpha_{\overline{\text{MS}}}(m_\tau^2)$ obtained from τ decays [1].

The determination of $\alpha_{\overline{\text{MS}}}$ at the Z^0 mass scale implies first to run up to the MS running mass for the bottom quark, m_b, with β-coefficients and $\Lambda_{\overline{\text{MS}}}$ estimated for 4 quark flavours, apply next the following matching formula [4],

$$\alpha_{\overline{\text{MS}}}^{N_f=5}(m_b) = \alpha_{\overline{\text{MS}}}^{N_f=4}(m_b)\left(1+\sum_n c_{n0}\left(\alpha_{\overline{\text{MS}}}^{N_f=4}(m_b)\right)^n\right) ,\tag{6}$$

where the coefficients c_{n0} can be found in ref. [22], and then run from the bottom mass up to the Z^0 mass scale. Thus, from our two estimates of $\Lambda_{\overline{\text{MS}}}$, we obtain: $\alpha_{\overline{\text{MS}}}(m_Z^2) = 0.1198(9)$ and $\alpha_{\overline{\text{MS}}}(m_Z^2) = 0.1203(11)$. Again, combining these two results and their errors added in quadrature, we will be left with

$$\alpha_{\overline{\text{MS}}}(m_Z^2) = 0.1200(14) \ , \tag{7}$$

also in good agreement with the current *world average* [4]: $\alpha_{\overline{\rm MS}}(m_Z^2) = 0.1184(7)$. It is worth to remark that this *world average* is strongly weighted by the HPQCD lattice (with 2+1 staggered fermions) result [5], $\alpha_{\overline{\rm MS}}(m_Z^2) = 0.1183(8)$, and by the one from non-singlet structure functions from DIS [23], $\alpha_{\overline{\rm MS}}(m_Z^2) = 0.1142(23)$, more than 2 σ's away from most of the other involved estimates. Without these two last estimates, the average would be 0.1197(12) and, if one just replaces the HPQCD result by the recent one form the PACS-CS lattice collaboration (with 2+1 Wilson improved fermions) [24], $\alpha_{\overline{\rm MS}}(m_Z^2) = 0.1205(8)(5)$, the average would be 0.1190(9), both in still better agreement with our estimate. Finally, the *world average* including the present result, with u, d, s and c realistic quarks in the loops, instead of the previous lattice one only with u, d and s, would be: $\alpha_{\overline{\rm MS}}(m_Z^2) = 0.1191(8)$.

It should be noted that we applied two different fitting strategies, taking different fitting windows and studying the impact of higher order OPE corrections, and no systematic effect have been observed. Our error analysis is based on the Jacknife method when we account for the fitted parameters, while the uncertainties on the lattice sizes (and the ratios of them) are properly propagated into the final estimates. Some systematic effect related to the use of the twisted-mass action for the dynamical quarks and to the lattice size determination at the chiral limit (not included in our budget error) could also appear and can be only excluded by the comparison with other lattice and experimental estimates.

CONCLUSIONS

We have presented the results for a first computation of the strong running coupling from lattice QCD simulations including u,d,s and c dynamical flavours. We applied the procedure of determining the ghost-gluon coupling renormalized in Taylor scheme over a large momenta window and then compare this with the running obtained in perturbative QCD and improved via non-perturbative OPE corrections. That procedure have been previously shown to work rather well when analysing lattice simulations with N_f=0 and 2 dynamical flavours and so happens here for N_f=2+1+1: our estimate for the strong running coupling at the τ-mass scale nicely agrees with those from τ-decays and, after being properly propagated down to the Z^0-mass scale, is pretty consistent with most of the estimates applied to obtain the current PDG *world average*, although slightly larger than the N_f=2+1 lattice result also used for this average.

Acknowledgements

We thank the IN2P3 (CNRS-Lyon), IDRIS (CNRS-Orsay) and apeNEXT (Rome) computing centers. This work has been partially supported by the Spanish MICINN research project FPA2011-23781 and by "Junta de Andalucia" P07FQM02962.

[1] S. Bethke, A. H. Hoang, S. Kluth, J. Schieck, I. W. Stewart, et al. (2011), * Temporary entry *, 1110.0016.
[2] Y. Aoki, PoS **LAT2009**, 012 (2009), 1005.2339.
[3] B. Blossier, P. Boucaud, M. Brinet, F. De Soto, X. Du, et al. (2011), * Temporary entry *, 1110.5829.
[4] K. Nakamura et al. (Particle Data Group), J. Phys. **G37**, 075021 (2010).
[5] C. Davies et al. (HPQCD Collaboration), Phys.Rev. **D78**, 114507 (2008), 0807.1687.
[6] R. Frezzotti, P. A. Grassi, S. Sint, and P. Weisz (Alpha), JHEP **08**, 058 (2001), hep-lat/0101001.
[7] R. Frezzotti and G. C. Rossi, Nucl. Phys. Proc. Suppl. **128**, 193 (2004), hep-lat/0311008.
[8] R. Baron, P. Boucaud, J. Carbonell, A. Deuzeman, V. Drach, et al. (ETM Collaboration), JHEP **1006**, 111 (2010), 1004.5284.
[9] R. Baron, et al. (ETM Collaboration), PoS **LATTICE2010**, 123 (2010), 1101.0518.
[10] P. Boucaud, F. De Soto, J. Leroy, A. Le Yaouanc, J. Micheli, O. Pène, and J. Rodríguez-Quintero, Phys.Rev. **D79**, 014508 (2009), 0811.2059.
[11] B. Blossier et al. (ETM), Phys. Rev. **D82**, 034510 (2010), 1005.5290.
[12] D. Becirevic, et al., Phys. Rev. **D60**, 094509 (1999), hep-ph/9903364; F. de Soto and C. Roiesnel, JHEP **0709**, 007 (2007), 0705.3523.
[13] J. A. Gracey, Phys. Lett. **B552**, 101 (2003), hep-th/0211144.
[14] K. Chetyrkin and A. Maier, JHEP **1001**, 092 (2010), 0911.0594.
[15] K. Chetyrkin and A. Retey (2000), hep-ph/0007088.
[16] P. Boucaud, A. Le Yaouanc, J. Leroy, J. Micheli, O. Pène, and J. Rodríguez-Quintero, Phys.Rev. **D63**, 114003 (2001), hep-ph/0101302; Phys.Lett. **B493**, 315 (2000), hep-ph/0008043.
[17] P. Boucaud et al., JHEP **01**, 046 (2002), hep-ph/0107278.
[18] P. Boucaud, J. Leroy, A. Le Yaouanc, A. Lokhov, J. Micheli, et al., JHEP **0601**, 037 (2006), hep-lat/0507005.
[19] F. V. Gubarev and V. I. Zakharov, Phys. Lett. **B501**, 28 (2001), hep-ph/0010096.
[20] D. Dudal, R. Sobreiro, S. Sorella, and H. Verschelde, Phys.Rev. **D72**, 014016 (2005), hep-th/0502183.
[21] E. Ruiz Arriola, P. O. Bowman, and W. Broniowski, Phys.Rev. **D70**, 097505 (2004), hep-ph/0408309.
[22] K. Chetyrkin, J. H. Kuhn, and C. Sturm, Nucl.Phys. **B744**, 121 (2006), hep-ph/0512060.
[23] J. Blumlein, H. Bottcher, and A. Guffanti, Nucl.Phys. **B774**, 182 (2007), hep-ph/0607200.
[24] S. Aoki et al. (PACS-CS), JHEP **10**, 053 (2009), 0906.3906.